



Unit 3

Disaster Medical Operations 1

Quiz 2

1. The first action to take when approaching a victim is:
 - ☐ a) Assess airway
 - ☐ b) Control bleeding
 - ☐ c) Check for pulse
 - ☐ d) Survey the area

 2. How many times will an attempt be made to open a victim's airway?
 - ☐ a) Once
 - ☐ b) Twice
 - ☐ c) Three times

 3. Signs and symptoms of shock include which of the following?
 - ☐ a) Pale skin
 - ☐ b) Cool skin temperature
 - ☐ c) Nausea
 - ☐ d) All of the above

 4. What is the treatment for a patient in shock?
 - ☐ a) Lay patient down
 - ☐ b) Elevate their feet 10-12 inches
 - ☐ c) Maintain an open airway
 - ☐ d) Maintain body temperature
 - ☐ e) All of the above

 5. Define Triage
-



Unit 3

Disaster Medical Operations 1

Quiz 2

6. What is the term for inadequate perfusion of the body's cells?

7. What methods are available to control bleeding in a disaster situation?

- ☐ a) Direct pressure
- ☐ b) Elevation
- ☐ c) Pressure points
- ☐ d) Tourniquet
- ☐ e) All of the above

8. All impaled objects should be left in place and stabilized with a bulky dressing unless the object is obstructing the patient's _____

9. The best method to open a victim's airway is to

- ☐ a) Massage their throat
- ☐ b) Blow in harder
- ☐ c) Turn person on their side
- ☐ d) Head tilt/chin lift

10. What does RPM stand for?

- ☐ a) Revolutions per minute
- ☐ b) Respiration, pulse, mental status
- ☐ c) Red, pink, magenta
- ☐ d) Rotate, push, mark